

## SOUTHERN COSMOLOGY SURVEY II: MASSIVE OPTICALLY-SELECTED CLUSTERS FROM 70 SQUARE DEGREES OF THE SZE COMMON SURVEY AREA

FELIPE MENANTEAU<sup>1</sup>, JOHN P. HUGHES<sup>1</sup>, L. FELIPE BARRIENTOS<sup>2</sup>, AMRUTA J. DESHPANDE<sup>1</sup>, MATT HILTON<sup>3</sup>,  
LEOPOLDO INFANTE<sup>2</sup>, RAUL JIMENEZ<sup>4</sup>, ARTHUR KOSOWSKY<sup>5</sup>, KAVILAN MOODLEY<sup>3</sup>, DAVID SPERGEL<sup>6</sup>, AND LICIA VERDE<sup>4</sup>

<sup>1</sup>Rutgers University, Department of Physics & Astronomy, 136 Frelinghuysen Rd, Piscataway, NJ 08854, USA

<sup>2</sup>Pontificia Universidad Católica de Chile, Departamento de Astronomía, Santiago, Chile

<sup>3</sup>University of KwaZulu-Natal, Astrophysics & Cosmology Research Unit, School of Mathematical Sciences, Durban, 4041, South Africa.

<sup>4</sup>ICREA & Institute for Sciences of the Cosmos (ICC), University of Barcelona, Martí i Franques 1, Barcelona 08034, Spain

<sup>5</sup>University of Pittsburgh, Physics & Astronomy Department, 100 Allen Hall, 3941 O'Hara Street, Pittsburgh, PA 15260, USA and

<sup>6</sup>Department of Astrophysical Sciences, Peyton Hall, Princeton University, Princeton, NJ 08544, USA

*ApJSS accepted*

### ABSTRACT

We present a catalog of 105 rich and massive ( $M > 3 \times 10^{14} M_{\odot}$ ) optically-selected clusters of galaxies extracted from 70 square-degrees of public archival *griz* imaging from the Blanco 4-m telescope acquired over 45 nights between 2005 and 2007. We use the clusters' optically-derived properties to estimate photometric redshifts, optical luminosities, richness, and masses. We complement the optical measurements with archival XMM-Newton and ROSAT X-ray data which provide additional luminosity and mass constraints on a modest fraction of the cluster sample. Two of our clusters show clear evidence for central lensing arcs; one of these has a spectacular large-diameter, nearly-complete Einstein Ring surrounding the brightest cluster galaxy. A strong motivation for this study is to identify the massive clusters that are expected to display prominent signals from the Sunyaev-Zeldovich Effect (SZE) and therefore be detected in the wide-area mm-band surveys being conducted by both the Atacama Cosmology Telescope and the South Pole Telescope. The optical sample presented here will be useful for verifying new SZE cluster candidates from these surveys, for testing the cluster selection function, and for stacking analyzes of the SZE data.

*Subject headings:* cosmic microwave background — cosmology: observations — galaxies: distances and redshifts — galaxies: clusters: general — large-scale structure of universe

### 1. INTRODUCTION

A new era of galaxy cluster surveys, based on measuring distortions in the cosmic microwave background (CMB), has begun. These distortions, known as the Sunyaev-Zel'dovich Effect (Sunyaev & Zel'dovich 1972), have been detected for the first time in untargeted surveys over large areas of the sky by two new mm-band experiments: the Atacama Cosmology Telescope (ACT) and the South Pole Telescope (SPT). First results from ACT (Hincks et al. 2009) and SPT (Staniszewski et al. 2009) offer a taste of the future potential these experiments hold for obtaining large samples of essentially mass-selected clusters to arbitrary redshifts and have also provided the first measurement of the microwave background at arc-minute angular scales (Lueker et al. 2009; Fowler et al. 2010).

Both ACT and SPT aim to provide unique samples of massive clusters of galaxies, selected by mass nearly independently of redshift, over a large area of the southern sky. While the number density of SZE-selected clusters can be used as a potentially strong probe of dark energy—as well as for studies of cluster physics, it is crucial to understand the systematics of SZE surveys by comparing with cluster identification using independent methods before the new cluster samples can be effectively used. For example the low amplitude of the SZ component in the high- $l$  CMB power spectrum (Lueker et al. 2009) or stacked clusters in WMAP (Komatsu et al. 2010) are recent issues that provide additional motivation for an independent search for clusters over the region

being surveyed in the SZ. Furthermore, although the new mm-band telescopes can be used to identify clusters, coordinated optical data are necessary for confirmation and to determine redshifts and other fundamental properties of the new clusters.

The last decade has seen significant effort to produce large and well-selected optical catalogs of cluster of galaxies that can be used in cosmological, large-scale structure and galaxy evolution studies. The first systematic attempts to generate large samples of clusters, and to define their richness, came from the Abell catalogs (Abell 1958; Abell et al. 1989) which searched for projected galaxy overdensities through visual inspection of photographic plates successfully identifying thousands of clusters. Although optical catalogs can be relatively inexpensive and efficient at detecting low mass systems, early attempts were known to suffer from significant projection effects along the line of sight. The advent of CCD cameras and the digitization of large photographic plates has enabled the development of new search algorithms for galaxy clusters using a combination of space, brightness and color information (i.e. photometric redshifts), minimizing projection issues (see Gal 2008, for a historical review of search methods). Among these algorithms are the pioneering implementation of a spatial matched filter technique (Postman et al. 1996) and its variants (Dong et al. 2008), the adaptive kernel technique (Gal et al. 2000, 2009), Voronoi tessellation (Ramella et al. 2001; Lopes et al. 2004) and methods exploiting the tight ridgeline in color-magnitude space of galaxies in clusters (Bower et al. 1992; Blakeslee et al. 2003) such as the Red

Cluster Sequence (RCS) (Gladders & Yee 2000) and the MaxBCG (Annis et al. 1999; Koester et al. 2007) techniques. This new wave of studies have produced large sets of well-defined optical cluster catalogs covering thousands of square-degrees (i.e. Goto et al. 2002; Gladders & Yee 2005; Koester et al. 2007; Gal et al. 2009) providing reliable richness-mass correlations (Becker et al. 2007; Johnston et al. 2007; Reyes et al. 2008; Sheldon et al. 2009) and establishing independent cosmological constraints (Rozo et al. 2010) using optical catalogs.


In this article we present new results from the Southern Cosmology Survey (SCS), our on-going multi-wavelength survey coordinated with ACT. Our first paper in this series, Menanteau et al. (2009), described our SCS imaging pipeline and presented a sample of new galaxy clusters from an 8 deg<sup>2</sup> optical imaging survey of the southern sky acquired at the Blanco 4-m telescope. Here, we complete our cluster analysis using all the 70 deg<sup>2</sup> contiguous imaging available, which represents a comoving volume of 0.076 Gpc<sup>3</sup> at  $z = 0.6$ . Throughout this paper we assume a flat cosmology with  $H_0 = 100h$  km s<sup>-1</sup> Mpc<sup>-1</sup>,  $h = 0.7$  and matter density  $\Omega_M = 0.3$ .

## 2. DATA SET AND ANALYSIS

Our cluster analysis is based on all of the public unprocessed imaging data available for download at the time of writing from the observations carried out for the Blanco Cosmology Survey<sup>1</sup> (BCS) proposal. This was a NOAO Large Survey project (05B-0043, PI: Joe Mohr) that was awarded 45 nights between 2005 and 2007 to carry out intermediate depth  $g, r, i, z$ -band observations on the Cerro Tololo InterAmerican Observatory (CTIO) Blanco 4-m telescope using the 8192 × 8192 pixel (0.36 deg<sup>2</sup>) MOSAIC-II camera. The survey originally aimed to target two square 50 deg<sup>2</sup> southern sky patches centered near declinations  $-55^\circ$  and  $-52^\circ$  at right ascensions of 23 hr and 5 hr respectively, that were to be contained in a larger SZE common region that both ACT and SPT would survey. After three years of observations the survey acquired 78 and 118 MOSAIC-II contiguous pointings at 23hr and 5hr respectively that correspond to 29 deg<sup>2</sup> and 41 deg<sup>2</sup> over each patch reaching a total of 70 deg<sup>2</sup> imaged in all *griz* filters. The pointing layout and the sky area covered for each patch is shown in Figure 1. The survey was awarded 15 extra nights at the end of 2008 to complete the proposed 100 deg<sup>2</sup>; our analysis does not include this extra year's worth of observations as the data are not publicly available yet.

### 2.1. Data Processing

The raw unprocessed images were downloaded from the National Virtual Observatory<sup>2</sup> portal and were processed following the same procedures as described in our initial analysis (Menanteau et al. 2009) where we provide a full description of the data analysis followed and associated data products. Here we briefly outline the steps involved and recent improvements from our original analysis. Our pipeline automatically handles the file associations and all of the initial standard CCD imaging tasks for each night's run as well as the secondary calibration steps (i.e., skyflats, fringe correction, cosmic-ray



**Figure 1.** The computed likelihood density map images centered at  $z = 0.3$  and width  $\Delta z = 0.1$  over the 23hr (top panel) 5hr (bottom panel) field. Dark regions in the image represent denser areas. In red we show the area covered by each of the 78 and 112 MOSAIC-II tiles at 23hr and 5hr respectively that comprise the area studied over three years of observations.


rejection, badpixel masks, world coordinate calibration, etc.) using a custom modified version of IRAF<sup>3</sup>/mscred (Valdes 1998). When observed, photometric standards from the Southern Hemisphere Standards Stars Catalog (Smith et al. 2007) were processed by the pipeline and a photometric zero-point was estimated in AB magnitudes. Hereafter all magnitude quoted are in the AB system. In table 1 we show the observing dates, photometric conditions and observed bands that comprise the dataset we analyzed and that were used for this paper. We note that the information that went into creating this table was extracted from the raw data and header information of the data files.

In general each MOSAIC II pointing consisted of exposures of  $2 \times 125s$ ,  $2 \times 300s$ ,  $3 \times 450s$  and  $3 \times 235s$  in the

<sup>1</sup> <http://cosmology.uiuc.edu/BCS/>

<sup>2</sup> <http://portal-nvo.noao.edu>

<sup>3</sup> IRAF is distributed by the National Optical Astronomy Observatory, which is operated by the AURA under cooperative agreement with NSF



**Figure 2.** The  $E(B - V)$  dust extinction maps created for the 5hr (left) and 23hr (right) fields in magnitudes using the Schlegel et al. (1998) maps of dust infrared emission. Both maps are showed with the same  $E(B - V)$  scale and size scales. Darker regions represent areas of higher Galactic dust absorption.

$g$ ,  $r$ ,  $i$  and  $z$ -bands respectively with offsets of 3 – 5 arcmin (within each filter). We adjusted the photometric zero points of non-photometric nights using the overlapping regions between tiles and their matched sources from adjacent photometric tiles providing a homogeneous photometric calibration across each region with typical variations below 0.03 mags. The 2007 season data at 23hr was particularly hard to match due to poor observing conditions and large variations within the individual exposures in each tile. We tackled this problem by first scaling all individual exposures comprising a tile to a common median scale; this step ensured that opposed edges of a combined frame could be effectively used to photometrically tie down adjacent tiles. This is a new feature of the pipeline.

All science images were mosaicked, aligned and median combined using SWarp (Bertin 2006) to a plate scale of  $0.''266/\text{pixel}$ . Source detection and photometry measurements for the science catalogs were performed using SExtractor (Bertin & Arnouts 1996). Large variations in the Galactic dust absorption on the 5hr field led us to implement dust correction for every source in each observed band in the science catalogs utilizing the infrared maps and C-routines provided by Schlegel et al. (1998). This is particularly important to obtain unbiased colors for photometric redshifts. In Figure 2 we show our custom generated  $E(B - V)$  Galactic dust absorption maps for the 5hr and 23hr regions, where darker regions represent areas of higher obscuration. From this figure we can see that in the 23hr region foreground Galactic absorption is negligible, while at 5hr it can be as high as  $E(B - V) \simeq 0.14$  magnitudes. Finally, we determine photometric redshifts from the four-band optical images and the Galactic-extinction-corrected magnitudes using BPZ (Benítez 2000) following the same procedure as discussed in Menanteau et al. (2009).

## 2.2. Finding Clusters and Defining Membership

In this section we describe our efforts to select clusters of galaxies from multi-wavelength optical imaging. To this objective we follow an identical procedure as the one we laid out in Menanteau et al. (2009) as part of our initial study of  $8 \text{ deg}^2$  using the 2005 observations at 23hr. We refer the reader there for an expanded and formal description of the method as well as for simulations that address the contamination and completeness of the search method. Here, we expanded our procedure to the full  $70 \text{ deg}^2$  of Blanco *griz* contiguous imaging. Our cluster search algorithm uses a matched filter approach based on the one described in Postman et al. (1996) to identify significant overdensities. We define membership and estimate richness for each peak in the overdensity maps using the MaxBCG prescription (Koester et al. 2007). Our method folds in the contributions from: a) a cluster spatial profile filter function, b) a luminosity weight and c) the BPZ redshift probability distribution from each source to generate likelihood density maps or a “filtered” galaxy catalog over the area covered by the survey as a function of redshift. We generate likelihood density maps with a constant pixel scale of 1.2 arcmin at  $\Delta z = 0.1$  intervals between  $0.1 < z < 0.8$  over the surveyed regions. In Figure 1 we show an example of a likelihood density map centered at  $z = 0.2$  on which we superpose outlines of the 78 and 112 tiles that define the 23hr and 5hr regions studied.

Because the mass of a cluster is not a direct observable, some observable proxy for mass needs to be used in order to obtain estimates for a given cluster sample. Such proxies include the X-ray flux and temperature (Reiprich & Böhringer 2002; Kravtsov et al. 2006; Rykoff et al. 2008b; Rozo et al. 2009a; Vikhlinin et al. 2009), weak lensing shear (Sheldon et al. 2009; Okabe et al. 2010) and optical galaxy richness of clusters (Becker et al. 2007; Rozo et al. 2009b). For this analysis we use the latest mass

**Table 1**  
2005-2007 Observations in the 23hr and 5hr Fields

| Date        | Photometric | # of Tiles Obs 05h |          |          |          | # of Tiles Obs 23h |          |          |          |
|-------------|-------------|--------------------|----------|----------|----------|--------------------|----------|----------|----------|
|             |             | <i>g</i>           | <i>r</i> | <i>i</i> | <i>z</i> | <i>g</i>           | <i>r</i> | <i>i</i> | <i>z</i> |
| 18 Nov 2005 | yes         | ...                | ...      | 1.7      | 1.3      | 1.0                | 1.0      | 3.0      | 3.0      |
| 19 Nov 2005 | yes         | ...                | ...      | 4.7      | 5.0      | 4.5                | 4.5      | 1.0      | 1.0      |
| 20 Nov 2005 | no          | ...                | ...      | 5.7      | 5.7      | 8.5                | 8.0      | ...      | ...      |
| 22 Nov 2005 | no          | ...                | ...      | ...      | ...      | ...                | ...      | ...      | ...      |
| 24 Nov 2005 | yes         | 9.0                | 9.0      | 1.7      | 1.7      | 5.5                | 5.5      | ...      | ...      |
| 26 Nov 2005 | yes         | 9.5                | 9.0      | 1.0      | 1.0      | 4.5                | 4.5      | 1.0      | 1.0      |
| 28 Nov 2005 | yes         | 4.0                | 3.0      | 4.0      | 4.0      | ...                | ...      | 2.7      | 2.7      |
| 30 Nov 2005 | yes         | 12.5               | 12.5     | ...      | ...      | 5.0                | 5.0      | 0.3      | 0.3      |
| 02 Dec 2005 | yes         | ...                | ...      | ...      | ...      | 3.5                | 3.5      | ...      | ...      |
| 04 Dec 2005 | no          | 9.0                | 9.0      | ...      | ...      | 2.0                | 2.0      | ...      | ...      |
| 05 Dec 2005 | no          | 2.5                | 2.0      | 2.7      | 2.7      | 2.0                | 2.0      | 1.3      | 1.7      |
| 06 Dec 2005 | yes         | ...                | ...      | 4.0      | 4.0      | ...                | ...      | 1.3      | 1.0      |
| 08 Dec 2005 | yes         | ...                | ...      | 4.0      | 4.0      | ...                | ...      | 2.3      | 2.7      |
| 10 Dec 2005 | yes         | ...                | ...      | 5.7      | 5.7      | ...                | ...      | 3.0      | 3.0      |
| 11 Dec 2005 | yes         | ...                | ...      | 4.3      | 4.3      | ...                | ...      | 3.0      | 3.3      |
| 23 Oct 2006 | yes         | 9.0                | 10.5     | ...      | ...      | ...                | ...      | 5.0      | 5.0      |
| 25 Oct 2006 | no          | ...                | ...      | ...      | ...      | ...                | ...      | ...      | ...      |
| 27 Oct 2006 | yes         | 10.5               | 7.0      | ...      | ...      | ...                | ...      | 5.0      | 5.3      |
| 28 Oct 2006 | no          | ...                | ...      | ...      | ...      | ...                | ...      | 5.3      | 5.0      |
| 29 Oct 2006 | yes         | 8.0                | 8.0      | 2.0      | 2.0      | ...                | ...      | 4.0      | 4.3      |
| 30 Oct 2006 | yes         | ...                | ...      | ...      | ...      | ...                | ...      | 5.0      | 5.3      |
| 31 Oct 2006 | yes         | ...                | ...      | 6.0      | 6.0      | ...                | ...      | 3.0      | 2.7      |
| 12 Dec 2006 | yes         | ...                | ...      | 3.0      | 1.7      | ...                | ...      | ...      | ...      |
| 13 Dec 2006 | yes         | ...                | ...      | 3.0      | 3.3      | 1.0                | 0.5      | ...      | ...      |
| 14 Dec 2006 | no          | ...                | ...      | 2.7      | 2.7      | ...                | ...      | ...      | ...      |
| 15 Dec 2006 | yes         | 4.0                | 4.0      | 5.7      | 5.7      | ...                | ...      | ...      | ...      |
| 16 Dec 2006 | no          | ...                | ...      | 4.0      | 3.7      | ...                | ...      | ...      | ...      |
| 17 Dec 2006 | yes         | 7.0                | 6.5      | 4.3      | 5.0      | ...                | ...      | ...      | ...      |
| 18 Dec 2006 | yes         | 8.0                | 8.5      | ...      | ...      | ...                | ...      | ...      | ...      |
| 19 Dec 2006 | yes         | 3.0                | 3.0      | 6.3      | 6.3      | ...                | ...      | ...      | ...      |
| 20 Dec 2006 | yes         | ...                | ...      | 4.0      | 4.0      | ...                | ...      | ...      | ...      |
| 21 Dec 2006 | yes         | ...                | ...      | 8.7      | 8.7      | ...                | ...      | ...      | ...      |
| 22 Dec 2006 | yes         | 3.5                | 3.5      | 2.3      | 2.0      | ...                | ...      | ...      | ...      |
| 23 Dec 2006 | yes         | 6.5                | 6.0      | 6.0      | 6.3      | ...                | ...      | ...      | ...      |
| 11 Sep 2007 | yes         | 2.5                | 3.0      | 2.3      | 2.7      | 4.5                | 5.0      | 3.3      | 4.0      |
| 12 Sep 2007 | yes         | ...                | ...      | ...      | ...      | 1.0                | 1.0      | 1.0      | 1.0      |
| 13 Sep 2007 | yes         | ...                | ...      | ...      | ...      | 1.5                | 1.0      | 1.3      | 1.3      |
| 14 Sep 2007 | yes         | 1.5                | 1.5      | 1.7      | 1.3      | 4.5                | 5.0      | 4.7      | 4.7      |
| 15 Sep 2007 | no          | 2.0                | 2.0      | 0.7      | 1.3      | 2.5                | 2.5      | 1.0      | 1.0      |
| 16 Sep 2007 | yes         | ...                | ...      | ...      | ...      | 2.5                | 2.5      | ...      | ...      |
| 17 Sep 2007 | yes         | 1.0                | ...      | ...      | ...      | ...                | ...      | 2.0      | 1.7      |
| 18 Sep 2007 | yes         | ...                | ...      | 2.3      | 2.0      | 3.0                | 2.5      | 7.0      | 6.7      |
| 19 Sep 2007 | yes         | ...                | ...      | ...      | ...      | ...                | ...      | 4.7      | 5.0      |
| 11 Oct 2007 | yes         | ...                | ...      | ...      | ...      | ...                | ...      | ...      | ...      |
| 12 Oct 2007 | no          | ...                | ...      | ...      | ...      | ...                | ...      | ...      | ...      |
| 30 Oct 2007 | no          | ...                | ...      | 1.0      | ...      | ...                | 0.5      | ...      | ...      |
| 31 Oct 2007 | no          | ...                | ...      | ...      | ...      | ...                | ...      | ...      | ...      |
| 01 Nov 2007 | yes         | ...                | ...      | 1.3      | 2.3      | 2.5                | 2.0      | ...      | ...      |
| 02 Nov 2007 | yes         | ...                | ...      | ...      | ...      | 5.0                | 5.0      | ...      | ...      |
| 03 Nov 2007 | yes         | ...                | ...      | 3.0      | 3.0      | 9.5                | 9.5      | ...      | ...      |
| 04 Nov 2007 | yes         | 5.0                | 5.5      | ...      | ...      | 3.0                | 3.0      | ...      | ...      |
| 06 Nov 2007 | yes         | ...                | ...      | ...      | ...      | ...                | ...      | 4.7      | 4.3      |
| 07 Nov 2007 | yes         | 1.0                | 2.0      | 0.7      | 0.7      | 1.0                | 1.0      | 1.7      | 2.0      |
| 08 Nov 2007 | yes         | ...                | ...      | ...      | ...      | ...                | ...      | ...      | ...      |

**Note.** — Observing conditions during the 2005-2007 runs of the BCS as well as the number of tiles observed on each night. We only show the dates in which tiles in the 23hr or 5hr contiguous regions were observed.

tracers for clusters of galaxies that are based on optically observed parameters (Johnston et al. 2007; Reyes et al. 2008) extracted from a sample of around 13,000 clusters from the Sloan Digital Sky Survey (SDSS) MaxBCG catalog (Koester et al. 2007). To use these mass tracers we define membership in similar fashion as Koester et al. (2007). We then visually inspected each candidate-cluster peak in the density maps and selected the brightest elliptical galaxy in the cluster (BCG), which was taken to be the center and initial redshift of the system. We then use galaxies photometrically classified as E or

E/S0s according to their BPZ spectral type and within a projected radius of  $0.5h^{-1}$  Mpc and redshift interval  $|z - z_o| = |\Delta z| = 0.05$  to obtain a local color-magnitude relation for each color combination as well as the cluster mean redshift,  $z_c$ , for all cluster members, using a  $3\sigma$  median sigma-clipping algorithm. We use these to determine  $N_{1\text{Mpc}} = N_{\text{gal}}$ , the number of galaxies within  $1h^{-1}\text{Mpc}$  of the cluster center. For our richness measurements we estimated the galaxy background contamination and implemented an appropriate background sub-

traction method following the same procedure described in Menanteau et al. (2009) (see section 3.1). We use a statistical removal of unrelated field galaxies with similar colors and redshifts that were projected along the line of sight to each cluster. We estimate the surface number density of ellipticals in an annulus surrounding the cluster (within  $R_{200} < r < 2R_{200}$ ) with  $\Delta z = 0.05$  and the same colors as the cluster members. We measure this background contribution around the outskirts of each cluster and obtain a corrected value  $N_{gal}$  which is used to compute  $R_{200}$  and then corresponding values of  $N_{200}^{gal}$  and  $L_{200}$ . The magnitude of the correction ranges between 15 – 20%. We will refer to the corrected values hereafter.

Finally we show, as examples of the depth and data quality, composite *gri* color images of several clusters in the 5hr and 23hr regions in Figures 3 and 4, respectively, that cover a wide range of redshifts.

### 3. RESULTS

#### 3.1. Optical Cluster Masses

The observational quantities required as input to the cluster mass scaling relation are  $N_{200}^{gal}$ ,  $L_{200}$ , and  $L_{BCG}$ . The cluster richness,  $N_{200}^{gal}$ , is the number of E and E/S0 galaxies within a given radius, originally defined by Hansen et al. (2005) as  $R_{200} = 0.156 N_{1Mpc}^{0.6} h^{-1}$  Mpc, with colors and luminosities that satisfy specific conditions for membership. Similarly,  $L_{200}$  is the total rest-frame integrated *r*-band luminosity, *k*-corrected to  $z = 0.25$ , of all member galaxies included in  $N_{200}^{gal}$ , and  $L_{BCG}$  is the similarly defined rest-frame *r*-band luminosity of the BCG. Reyes et al. (2008) provide power-law functions for both the luminosity-mass and richness-mass relations (see section 5.2.1 in their paper).

Both Johnston et al. (2007) and Reyes et al. (2008) found that the luminosity-mass and richness-mass relations were well described by power-law functions and they measured the normalizations and slopes in these relations using  $\chi^2$  minimization. We computed the two fitting functions based on  $L_{200}$  and  $N_{200}^{gal}$ , (see section 5.2.1 from Reyes et al. 2008 for full details), which are described as:

$$M(N_{200}, L_{BCG}) = M_N^0 (N_{200}/20)^{\alpha_N} (L_{BCG}/\bar{L}_{BCG}^{(N)})^{\gamma_N} \quad (1)$$

and

$$M(L_{200}, L_{BCG}) = M_L^0 (L_{200}/40)^{\alpha_L} (L_{BCG}/\bar{L}_{BCG}^{(L)})^{\gamma_L} \quad (2)$$

where  $M$  is the mass observational equivalent of  $M_{200\bar{\rho}}$  (i.e. the halo mass enclosed within a radius of spherical volume within which the mean density is 200 times the average density) in units of  $10^{14} M_\odot$ ,  $L_{200}$  is in units of  $10^{10} h^{-2} L_\odot$  and the  $L_{BCG}$  dependence is normalized by its mean value. This is also described by a power-law function for a given value of  $L_{200}$  and  $N_{200}^{gal}$ :

$$\bar{L}_{BCG}^{(N)} \equiv \bar{L}_{BCG}(N_{200}) = a_N N_{200}^{b_N} \quad (3)$$

and

$$\bar{L}_{BCG}^{(L)} \equiv \bar{L}_{BCG}(L_{200}) = a_L L_{200}^{b_L} \quad (4)$$

The published best-fitting parameters for  $M^0$ ,  $\alpha$  and  $\gamma$  in Eqs. (1) and (2) as well as the new erratum-corrected

values of  $a$  and  $b$  (R. Reyes, private communication) for Eqs. (3) and (4) are shown in Table 2. These recent changes in the values of the  $a$  and  $b$  parameters in Reyes et al. (2008) have implications for our mass estimation. Specifically, the changes translate into a decrease in mass when compared to our previous analysis of Menanteau et al. (2009) which used the initial parameters from their pre-print paper. As we discuss below, this change also affects the recovery of clusters in Menanteau et al. (2009) for the sky region that overlaps with this study.

We use this parametrization to obtain the optical mass estimates  $M(N_{200})$  and  $M(L_{200})$ , based on  $N_{200}^{gal}$  and  $L_{200}$ , respectively, for all of the visually inspected clusters. Based on these estimates we defined our catalog of massive clusters by selecting all systems with either  $M(N_{200}) > 3 \times 10^{14} M_\odot$  or  $M(L_{200}) > 3 \times 10^{14} M_\odot$  and  $N_{gal} > 10$ . This defines the sample: 61 systems in the 5hr region and 44 in the 23hr region for a total of 105 clusters over 70 deg<sup>2</sup> of multi-band imaging. This mass threshold is aimed to include the upcoming  $z < 0.8$  significant SZE detections from SPT and ACT. In Tables 3 and 4 we display the optical properties, photometric redshifts, positions and mass estimates for all clusters at 23hr and 5hr respectively. In these tables we provide for each cluster the photometric redshift of the BCG as well as the mean photometric redshift for the system which was estimated using galaxies within 250 kpc of the BCG.

As we discussed in Menanteau & Hughes (2009) the expected luminosity range of BCGs in rich clusters has been already observationally established from the SDSS (Loh & Strauss 2006; Koester et al. 2007) and it can be used as an additional probe to confirm the presence of a massive cluster. To this end in Figure 5 we compare the observed *r*-band magnitudes of all BCGs in our sample as a function of redshift to a parametrization of the observed *r*-band of SDSS BCGs (dashed curve). This corresponds to the  $M^* - 1.5$  prescription from Loh & Strauss (2006) where we have used  $M^*$  from Blanton et al. (2003) and allowed it to evolve passively with redshift. We see in the figure that all of the sources lie very close or well below (i.e., are intrinsically brighter) than the model curve as we would expect for the BCG in a rich cluster.


In our previous study (Menanteau et al. 2009) we presented a similar cluster analysis and mass estimation over 8 deg<sup>2</sup> area in the 23hr region which is fully contained in the current 70 deg<sup>2</sup> dataset. However, due to the changes in the mass parametrization prescription we did not recover all the massive clusters from Menanteau et al. (2009) even though the nominal mass threshold was the same ( $M > 3 \times 10^{14} M_\odot$ ) as here. Therefore, only 3 out of 8 clusters from Table 5 of Menanteau et al. (2009) are massive enough to appear in the current paper's Table 3, while the remaining clusters, although detected, now have masses that fall below the  $3 \times 10^{14} M_\odot$  mass limit. Similarly, when comparing with the weak lensing mass estimates of McInnes et al. (2009) we can only match the same 3 sources with clusters in their analysis (see Table 2 of their paper). On the other hand, there is now better agreement between the updated  $M(L_{200})$  values and the weak lensing masses for the matched clusters.






**Figure 3.** Composite *gri* color image for nine newly discovered massive SCS clusters in the 5hr field. We indicate the cluster's redshift using the median value from galaxy members within 250 kpc of the cluster's center.





**Figure 4.** Composite *gri* color image for nine newly discovered massive SCS clusters in the 23hr field. We indicate the cluster's redshift using the median value from galaxy members within 250 kpc of the cluster's center.



**Figure 5.** The *r*-band observed magnitude for the BCGs in our sample as a function of the redshift of the cluster (filled circles). We also show the  $M^* - 1.5$  BCG-redshift relationship from Loh & Strauss (2006) as the dashed curve.

### 3.2. Lensing Rate

Out of the 105 massive clusters in the sample we report two systems (see the two lower panels of Figure 4) with obvious evidence of arcs near the BCG. Of these two lensing clusters, SCSO 235138–545253 at  $z = 0.33$  is a potentially unique system in that it shows a large diameter ( $\sim 15''$ ), nearly complete Einstein Ring embedded in a massive cluster. Although our density of lensing clusters is not widely different from the Gladders et al. (2003) RCS sample of 5 lensing clusters over  $\sim 90 \text{ deg}^2$ , all of the Gladders et al. clusters are at  $z > 0.64$ , while both of our systems are at  $z < 0.54$  making them completely exclusive in redshift. Moreover, none of the Gladders et al. (2003) sample shows anything close to a complete Einstein Ring.

### 3.3. Correlation with Known Sources

We queried the NASA/IPCA Extragalactic Database (NED)<sup>4</sup> for catalogued clusters from *ROSAT*, Abell et al. (1989), ACT (Hincks et al. 2009) and SPT (Staniszewski et al. 2009; Menanteau & Hughes 2009) within a  $3'$  radius of the location of each SCS cluster (see Table 5). In some cases there was a catalogued galaxy from the 2 Micron All Sky Survey (Skrutskie et al. 2006, NED ID: 2MASX), which we report if it is within  $10''$  of the BCG. We also take note of radio sources within  $1'$  of our cluster positions since these could potentially bias the cluster SZ signal. We only found radio sources from the Sydney University Molonglo Sky Survey (SUMSS) at 843 MHz (Mauch et al. 2003). Finally we find one unidentified bright X-ray source from the *ROSAT* All Sky Survey (RASS) (Voges et al. 1999) that is coincident with one of our clusters.

It is interesting to note that although we recover two of the SZE-selected clusters from the first SPT results (Staniszewski et al. 2009), there are two others from

<sup>4</sup> <http://nedwww.ipac.caltech.edu/>

**Table 2**  
Mass-richness power-law function Best fitting parameters from Reyes et al. (2008)

| Redshift          | $(10^{10} h^{-2} L_{\odot})$ |       | $b_N$ | $b_L$ | $M_N^0$         | $\alpha_N$      | $\gamma_N$      | $M_L^0$         | $\alpha_L$      | $\gamma_L$      |
|-------------------|------------------------------|-------|-------|-------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|
|                   | $a_N$                        | $a_L$ |       |       |                 |                 |                 |                 |                 |                 |
| $0.10 < z < 0.23$ | 1.54                         | 0.61  | 0.41  | 0.67  | $1.27 \pm 0.08$ | $1.20 \pm 0.09$ | $0.71 \pm 0.14$ | $1.81 \pm 0.15$ | $1.27 \pm 0.17$ | $0.40 \pm 0.23$ |
| $0.23 < z < 0.70$ | 1.64                         | 0.58  | 0.43  | 0.66  | $1.57 \pm 0.14$ | $1.12 \pm 0.15$ | $0.34 \pm 0.24$ | $1.76 \pm 0.22$ | $1.30 \pm 0.29$ | $0.26 \pm 0.41$ |

**Table 3**  
Optical Clusters with  $M(N_{200}) > 3 \times 10^{14} M_{\odot}$  or  $M(L_{200}) > 3 \times 10^{14} M_{\odot}$  in the 23hr field

| ID                  | $z_{\text{cluster}}$ | $z_{\text{BCG}}$ | $N_{\text{gal}}$ | $N_{200}^{\text{gal}}$ | $L_{200}[L_{\odot}]$                        | $M(N_{200})$         | $M(L_{200})$<br>( $M_{\odot}$ ) |
|---------------------|----------------------|------------------|------------------|------------------------|---------------------------------------------|----------------------|---------------------------------|
| SCSO J231340–521919 | 0.21                 | 0.21             | $61.2 \pm 7.8$   | $124.8 \pm 11.2$       | $2.0 \times 10^{12} \pm 2.3 \times 10^{11}$ | $3.1 \times 10^{14}$ | $2.9 \times 10^{14}$            |
| SCSO J231403–564710 | 0.60                 | 0.60             | $22.7 \pm 4.8$   | $22.4 \pm 4.7$         | $1.2 \times 10^{12} \pm 5.2 \times 10^{10}$ | $2.2 \times 10^{14}$ | $3.5 \times 10^{14}$            |
| SCSO J231455–555308 | 0.21                 | 0.23             | $59.6 \pm 7.7$   | $79.9 \pm 8.9$         | $2.0 \times 10^{12} \pm 2.2 \times 10^{11}$ | $4.6 \times 10^{14}$ | $4.7 \times 10^{14}$            |
| SCSO J231511–523322 | 0.36                 | 0.39             | $29.0 \pm 5.4$   | $33.6 \pm 5.8$         | $2.3 \times 10^{12} \pm 1.0 \times 10^{11}$ | $3.9 \times 10^{14}$ | $8.2 \times 10^{14}$            |
| SCSO J231629–554535 | 0.51                 | 0.53             | $33.9 \pm 5.8$   | $38.1 \pm 6.2$         | $2.8 \times 10^{12} \pm 1.1 \times 10^{11}$ | $5.2 \times 10^{14}$ | $1.1 \times 10^{15}$            |
| SCSO J231651–545356 | 0.36                 | 0.36             | $35.5 \pm 6.0$   | $40.3 \pm 6.3$         | $9.6 \times 10^{11} \pm 9.6 \times 10^{10}$ | $3.9 \times 10^{14}$ | $2.7 \times 10^{14}$            |
| SCSO J231717–565723 | 0.74                 | 0.73             | $21.0 \pm 4.6$   | $21.8 \pm 4.7$         | $1.4 \times 10^{12} \pm 5.2 \times 10^{10}$ | $1.7 \times 10^{14}$ | $3.3 \times 10^{14}$            |
| SCSO J231848–561711 | 0.51                 | 0.51             | $28.8 \pm 5.4$   | $31.4 \pm 5.6$         | $1.6 \times 10^{12} \pm 9.4 \times 10^{10}$ | $3.4 \times 10^{14}$ | $5.2 \times 10^{14}$            |
| SCSO J231930–563858 | 0.36                 | 0.37             | $35.4 \pm 6.0$   | $38.4 \pm 6.2$         | $1.1 \times 10^{12} \pm 8.2 \times 10^{10}$ | $3.5 \times 10^{14}$ | $2.9 \times 10^{14}$            |
| SCSO J232001–565222 | 0.80                 | 0.80             | $18.0 \pm 4.2$   | $13.8 \pm 3.7$         | $1.3 \times 10^{12} \pm 5.4 \times 10^{10}$ | $1.8 \times 10^{14}$ | $4.6 \times 10^{14}$            |
| SCSO J232423–565705 | 0.75                 | 0.75             | $25.0 \pm 5.0$   | $30.6 \pm 5.5$         | $2.5 \times 10^{12} \pm 8.0 \times 10^{10}$ | $2.3 \times 10^{14}$ | $6.5 \times 10^{14}$            |
| SCSO J232437–530047 | 0.73                 | 0.75             | $38.7 \pm 6.2$   | $63.9 \pm 8.0$         | $2.2 \times 10^{12} \pm 2.1 \times 10^{11}$ | $3.9 \times 10^{14}$ | $4.7 \times 10^{14}$            |
| SCSO J232529–532420 | 0.74                 | 0.71             | $40.8 \pm 6.4$   | $46.5 \pm 6.8$         | $2.2 \times 10^{12} \pm 2.3 \times 10^{11}$ | $2.6 \times 10^{14}$ | $4.5 \times 10^{14}$            |
| SCSO J232612–531858 | 0.15                 | 0.13             | $77.2 \pm 8.8$   | $69.4 \pm 8.3$         | $1.8 \times 10^{12} \pm 3.2 \times 10^{11}$ | $2.5 \times 10^{14}$ | $3.1 \times 10^{14}$            |
| SCSO J232627–531512 | 0.74                 | 0.74             | $39.2 \pm 6.3$   | $42.0 \pm 6.5$         | $2.5 \times 10^{12} \pm 2.3 \times 10^{11}$ | $3.2 \times 10^{14}$ | $6.6 \times 10^{14}$            |
| SCSO J232637–533911 | 0.76                 | 0.77             | $34.4 \pm 5.9$   | $56.1 \pm 7.5$         | $3.1 \times 10^{12} \pm 2.2 \times 10^{11}$ | $4.5 \times 10^{14}$ | $8.6 \times 10^{14}$            |
| SCSO J232653–524149 | 0.11                 | 0.11             | $52.8 \pm 7.3$   | $87.1 \pm 9.3$         | $1.2 \times 10^{12} \pm 1.8 \times 10^{11}$ | $3.1 \times 10^{14}$ | $2.2 \times 10^{14}$            |
| SCSO J232724–533553 | 0.74                 | 0.74             | $56.6 \pm 7.5$   | $103.1 \pm 10.2$       | $6.0 \times 10^{12} \pm 5.4 \times 10^{11}$ | $9.2 \times 10^{14}$ | $2.0 \times 10^{15}$            |
| SCSO J232800–535152 | 0.74                 | 0.75             | $36.8 \pm 6.1$   | $63.9 \pm 8.0$         | $1.4 \times 10^{12} \pm 2.0 \times 10^{11}$ | $3.7 \times 10^{14}$ | $2.8 \times 10^{14}$            |
| SCSO J232811–533847 | 0.74                 | 0.74             | $43.1 \pm 6.6$   | $47.2 \pm 6.9$         | $2.1 \times 10^{12} \pm 2.3 \times 10^{11}$ | $3.4 \times 10^{14}$ | $5.2 \times 10^{14}$            |
| SCSO J232825–522814 | 0.73                 | 0.70             | $35.0 \pm 5.9$   | $47.3 \pm 6.9$         | $2.9 \times 10^{12} \pm 1.4 \times 10^{11}$ | $4.3 \times 10^{14}$ | $9.0 \times 10^{14}$            |
| SCSO J232827–531414 | 0.36                 | 0.35             | $34.7 \pm 5.9$   | $39.5 \pm 6.3$         | $1.5 \times 10^{12} \pm 1.1 \times 10^{11}$ | $3.0 \times 10^{14}$ | $3.7 \times 10^{14}$            |
| SCSO J232856–552428 | 0.57                 | 0.57             | $20.3 \pm 4.5$   | $19.6 \pm 4.4$         | $1.2 \times 10^{12} \pm 5.2 \times 10^{10}$ | $2.0 \times 10^{14}$ | $3.5 \times 10^{14}$            |
| SCSO J232916–522910 | 0.73                 | 0.74             | $39.8 \pm 6.3$   | $58.1 \pm 7.6$         | $3.3 \times 10^{12} \pm 1.6 \times 10^{11}$ | $4.5 \times 10^{14}$ | $9.2 \times 10^{14}$            |
| SCSO J233006–524035 | 0.73                 | 0.71             | $36.4 \pm 6.0$   | $43.8 \pm 6.6$         | $2.0 \times 10^{12} \pm 1.4 \times 10^{11}$ | $3.2 \times 10^{14}$ | $5.0 \times 10^{14}$            |
| SCSO J233227–535827 | 0.35                 | 0.32             | $41.0 \pm 6.4$   | $42.4 \pm 6.5$         | $9.3 \times 10^{11} \pm 9.0 \times 10^{10}$ | $3.7 \times 10^{14}$ | $2.4 \times 10^{14}$            |
| SCSO J233232–522016 | 0.36                 | 0.37             | $37.4 \pm 6.1$   | $43.2 \pm 6.6$         | $8.9 \times 10^{11} \pm 8.2 \times 10^{10}$ | $4.0 \times 10^{14}$ | $2.4 \times 10^{14}$            |
| SCSO J233329–521513 | 0.51                 | 0.50             | $36.8 \pm 6.1$   | $42.2 \pm 6.5$         | $1.7 \times 10^{12} \pm 1.0 \times 10^{11}$ | $3.6 \times 10^{14}$ | $4.6 \times 10^{14}$            |
| SCSO J233420–542732 | 0.56                 | 0.55             | $31.7 \pm 5.6$   | $37.5 \pm 6.1$         | $1.5 \times 10^{12} \pm 9.8 \times 10^{10}$ | $2.7 \times 10^{14}$ | $3.6 \times 10^{14}$            |
| SCSO J233544–535115 | 0.51                 | 0.51             | $58.8 \pm 7.7$   | $96.8 \pm 9.8$         | $4.3 \times 10^{12} \pm 2.1 \times 10^{11}$ | $8.1 \times 10^{14}$ | $1.3 \times 10^{15}$            |
| SCSO J233556–560602 | 0.64                 | 0.63             | $14.2 \pm 3.8$   | $19.8 \pm 4.5$         | $1.3 \times 10^{12} \pm 7.3 \times 10^{10}$ | $2.5 \times 10^{14}$ | $4.7 \times 10^{14}$            |
| SCSO J233607–535235 | 0.54                 | 0.53             | $60.1 \pm 7.8$   | $83.3 \pm 9.1$         | $5.3 \times 10^{12} \pm 2.6 \times 10^{11}$ | $1.0 \times 10^{15}$ | $2.2 \times 10^{15}$            |
| SCSO J233726–565655 | 0.50                 | 0.52             | $26.7 \pm 5.2$   | $31.1 \pm 5.6$         | $1.2 \times 10^{12} \pm 7.6 \times 10^{10}$ | $2.7 \times 10^{14}$ | $3.0 \times 10^{14}$            |
| SCSO J233807–531223 | 0.47                 | 0.49             | $46.1 \pm 6.8$   | $53.3 \pm 7.3$         | $2.1 \times 10^{12} \pm 1.8 \times 10^{11}$ | $4.8 \times 10^{14}$ | $6.2 \times 10^{14}$            |
| SCSO J233931–544525 | 0.73                 | 0.71             | $29.6 \pm 5.4$   | $33.6 \pm 5.8$         | $1.6 \times 10^{12} \pm 1.1 \times 10^{11}$ | $2.7 \times 10^{14}$ | $4.1 \times 10^{14}$            |
| SCSO J234012–541907 | 0.59                 | 0.62             | $25.9 \pm 5.1$   | $26.2 \pm 5.1$         | $1.5 \times 10^{12} \pm 9.8 \times 10^{10}$ | $2.6 \times 10^{14}$ | $4.4 \times 10^{14}$            |
| SCSO J234138–545210 | 0.55                 | 0.56             | $25.7 \pm 5.1$   | $26.3 \pm 5.1$         | $1.1 \times 10^{12} \pm 7.8 \times 10^{10}$ | $2.8 \times 10^{14}$ | $3.5 \times 10^{14}$            |
| SCSO J234156–530848 | 0.49                 | 0.49             | $44.7 \pm 6.7$   | $77.1 \pm 8.8$         | $2.7 \times 10^{12} \pm 1.7 \times 10^{11}$ | $7.8 \times 10^{14}$ | $8.9 \times 10^{14}$            |
| SCSO J234703–535051 | 0.56                 | 0.55             | $21.6 \pm 4.7$   | $22.0 \pm 4.7$         | $1.1 \times 10^{12} \pm 5.6 \times 10^{10}$ | $2.0 \times 10^{14}$ | $3.0 \times 10^{14}$            |
| SCSO J234917–545521 | 0.73                 | 0.72             | $17.9 \pm 4.2$   | $17.4 \pm 4.2$         | $1.1 \times 10^{12} \pm 6.7 \times 10^{10}$ | $1.8 \times 10^{14}$ | $3.2 \times 10^{14}$            |
| SCSO J235055–530124 | 0.46                 | 0.48             | $41.4 \pm 6.4$   | $56.3 \pm 7.5$         | $1.4 \times 10^{12} \pm 1.1 \times 10^{11}$ | $4.8 \times 10^{14}$ | $3.6 \times 10^{14}$            |
| SCSO J235138–545253 | 0.33                 | 0.31             | $71.4 \pm 8.4$   | $122.3 \pm 11.1$       | $2.5 \times 10^{12} \pm 2.3 \times 10^{11}$ | $1.1 \times 10^{15}$ | $7.4 \times 10^{14}$            |
| SCSO J235233–564348 | 0.74                 | 0.72             | $17.1 \pm 4.1$   | $16.2 \pm 4.0$         | $1.1 \times 10^{12} \pm 5.2 \times 10^{10}$ | $1.7 \times 10^{14}$ | $3.1 \times 10^{14}$            |
| SCSO J235454–563311 | 0.51                 | 0.50             | $33.1 \pm 5.8$   | $39.4 \pm 6.3$         | $1.7 \times 10^{12} \pm 8.0 \times 10^{10}$ | $3.8 \times 10^{14}$ | $5.0 \times 10^{14}$            |

**Note.** — Catalog with the optical properties of clusters with mass estimates  $> 3 \times 10^{14} M_{\odot}$  in the 23hr region. For each cluster we note the BCGs photometric redshift and the median photometric redshift for the clusters using the members within 250 kpc of the center of the cluster. The ID is based on the position of the BCG.



**Table 4**  
Optical Clusters with  $M(N_{200}) > 3 \times 10^{14} M_{\odot}$  or  $M(L_{200}) > 3 \times 10^{14} M_{\odot}$  in the 5hr field

| ID                  | $z_{\text{cluster}}$ | $z_{\text{BCG}}$ | $N_{\text{gal}}$ | $N_{200}^{\text{gal}}$ | $L_{200}[L_{\odot}]$                        | $M(N_{200})$<br>( $M_{\odot}$ ) | $M(L_{200})$<br>( $M_{\odot}$ ) |
|---------------------|----------------------|------------------|------------------|------------------------|---------------------------------------------|---------------------------------|---------------------------------|
| SCSO J050854–513048 | 0.70                 | 0.70             | $25.0 \pm 5.0$   | $23.9 \pm 4.9$         | $1.3 \times 10^{12} \pm 1.4 \times 10^{11}$ | $2.4 \times 10^{14}$            | $3.8 \times 10^{14}$            |
| SCSO J050857–535837 | 0.76                 | 0.78             | $37.0 \pm 6.1$   | $48.5 \pm 7.0$         | $3.5 \times 10^{12} \pm 2.4 \times 10^{11}$ | $5.0 \times 10^{14}$            | $1.2 \times 10^{15}$            |
| SCSO J050902–520704 | 0.58                 | 0.58             | $35.3 \pm 5.9$   | $37.0 \pm 6.1$         | $1.9 \times 10^{12} \pm 1.7 \times 10^{11}$ | $2.8 \times 10^{14}$            | $4.8 \times 10^{14}$            |
| SCSO J050926–522227 | 0.67                 | 0.70             | $30.9 \pm 5.6$   | $33.5 \pm 5.8$         | $1.6 \times 10^{12} \pm 1.8 \times 10^{11}$ | $2.5 \times 10^{14}$            | $4.0 \times 10^{14}$            |
| SCSO J051023–544455 | 0.39                 | 0.37             | $45.2 \pm 6.7$   | $51.0 \pm 7.1$         | $1.1 \times 10^{12} \pm 1.7 \times 10^{11}$ | $3.5 \times 10^{14}$            | $2.3 \times 10^{14}$            |
| SCSO J051112–523112 | 0.73                 | 0.73             | $31.7 \pm 5.6$   | $34.4 \pm 5.9$         | $2.2 \times 10^{12} \pm 2.2 \times 10^{11}$ | $3.3 \times 10^{14}$            | $6.6 \times 10^{14}$            |
| SCSO J051136–561045 | 0.70                 | 0.71             | $30.9 \pm 5.6$   | $35.0 \pm 5.9$         | $2.9 \times 10^{12} \pm 1.6 \times 10^{11}$ | $3.2 \times 10^{14}$            | $8.7 \times 10^{14}$            |
| SCSO J051144–511416 | 0.48                 | 0.48             | $33.3 \pm 5.8$   | $40.1 \pm 6.3$         | $1.6 \times 10^{12} \pm 1.8 \times 10^{11}$ | $2.9 \times 10^{14}$            | $3.7 \times 10^{14}$            |
| SCSO J051145–515430 | 0.70                 | 0.70             | $32.7 \pm 5.7$   | $32.4 \pm 5.7$         | $1.6 \times 10^{12} \pm 2.1 \times 10^{11}$ | $2.2 \times 10^{14}$            | $3.5 \times 10^{14}$            |
| SCSO J051207–514204 | 0.48                 | 0.46             | $37.7 \pm 6.1$   | $49.1 \pm 7.0$         | $1.6 \times 10^{12} \pm 1.9 \times 10^{11}$ | $3.5 \times 10^{14}$            | $3.9 \times 10^{14}$            |
| SCSO J051225–505913 | 0.70                 | 0.70             | $49.8 \pm 7.0$   | $54.6 \pm 7.4$         | $3.0 \times 10^{12} \pm 3.2 \times 10^{11}$ | $4.9 \times 10^{14}$            | $9.2 \times 10^{14}$            |
| SCSO J051240–513941 | 0.66                 | 0.66             | $33.8 \pm 5.8$   | $30.8 \pm 5.5$         | $1.5 \times 10^{12} \pm 1.9 \times 10^{11}$ | $2.2 \times 10^{14}$            | $3.5 \times 10^{14}$            |
| SCSO J051245–502028 | 0.62                 | 0.62             | $24.2 \pm 4.9$   | $23.3 \pm 4.8$         | $2.6 \times 10^{12} \pm 5.4 \times 10^{10}$ | $2.8 \times 10^{14}$            | $9.4 \times 10^{14}$            |
| SCSO J051258–542153 | 0.67                 | 0.68             | $18.5 \pm 4.3$   | $18.4 \pm 4.3$         | $1.3 \times 10^{12} \pm 1.2 \times 10^{11}$ | $2.0 \times 10^{14}$            | $4.0 \times 10^{14}$            |
| SCSO J051400–512635 | 0.72                 | 0.73             | $54.7 \pm 7.4$   | $81.7 \pm 9.0$         | $5.7 \times 10^{12} \pm 4.9 \times 10^{11}$ | $7.0 \times 10^{14}$            | $1.8 \times 10^{15}$            |
| SCSO J051412–514004 | 0.67                 | 0.66             | $44.8 \pm 6.7$   | $60.9 \pm 7.8$         | $2.5 \times 10^{12} \pm 3.1 \times 10^{11}$ | $4.2 \times 10^{14}$            | $6.0 \times 10^{14}$            |
| SCSO J051457–514345 | 0.69                 | 0.69             | $36.4 \pm 6.0$   | $37.4 \pm 6.1$         | $4.0 \times 10^{12} \pm 3.4 \times 10^{11}$ | $4.1 \times 10^{14}$            | $1.5 \times 10^{15}$            |
| SCSO J051542–514017 | 0.73                 | 0.73             | $40.4 \pm 6.3$   | $53.4 \pm 7.3$         | $2.4 \times 10^{12} \pm 3.3 \times 10^{11}$ | $4.0 \times 10^{14}$            | $6.3 \times 10^{14}$            |
| SCSO J051558–543906 | 0.66                 | 0.64             | $33.7 \pm 5.8$   | $38.4 \pm 6.2$         | $1.5 \times 10^{12} \pm 1.8 \times 10^{11}$ | $2.8 \times 10^{14}$            | $3.5 \times 10^{14}$            |
| SCSO J051613–542620 | 0.38                 | 0.36             | $45.9 \pm 6.8$   | $65.2 \pm 8.1$         | $4.3 \times 10^{12} \pm 2.4 \times 10^{11}$ | $6.8 \times 10^{14}$            | $1.5 \times 10^{15}$            |
| SCSO J051637–543001 | 0.23                 | 0.25             | $127.0 \pm 11.3$ | $180.3 \pm 13.4$       | $5.3 \times 10^{12} \pm 1.0 \times 10^{12}$ | $1.8 \times 10^{15}$            | $1.9 \times 10^{15}$            |
| SCSO J051755–555727 | 0.66                 | 0.66             | $20.4 \pm 4.5$   | $18.2 \pm 4.3$         | $1.2 \times 10^{12} \pm 1.1 \times 10^{11}$ | $1.7 \times 10^{14}$            | $3.4 \times 10^{14}$            |
| SCSO J051933–554243 | 0.69                 | 0.70             | $25.0 \pm 5.0$   | $23.8 \pm 4.9$         | $2.5 \times 10^{12} \pm 1.8 \times 10^{11}$ | $2.9 \times 10^{14}$            | $9.1 \times 10^{14}$            |
| SCSO J051935–554916 | 0.75                 | 0.75             | $25.8 \pm 5.1$   | $25.7 \pm 5.1$         | $1.6 \times 10^{12} \pm 1.7 \times 10^{11}$ | $2.2 \times 10^{14}$            | $4.3 \times 10^{14}$            |
| SCSO J052051–561804 | 0.74                 | 0.73             | $55.4 \pm 7.5$   | $91.0 \pm 9.5$         | $4.5 \times 10^{12} \pm 3.7 \times 10^{11}$ | $7.1 \times 10^{14}$            | $1.3 \times 10^{15}$            |
| SCSO J052113–510418 | 0.64                 | 0.61             | $58.2 \pm 7.6$   | $85.1 \pm 9.2$         | $4.8 \times 10^{12} \pm 4.9 \times 10^{11}$ | $7.9 \times 10^{14}$            | $1.6 \times 10^{15}$            |
| SCSO J052200–502700 | 0.50                 | 0.47             | $74.0 \pm 8.6$   | $133.7 \pm 11.6$       | $4.1 \times 10^{12} \pm 4.0 \times 10^{11}$ | $9.4 \times 10^{14}$            | $1.1 \times 10^{15}$            |
| SCSO J052533–551818 | 0.72                 | 0.72             | $19.1 \pm 4.4$   | $15.3 \pm 3.9$         | $1.6 \times 10^{12} \pm 1.4 \times 10^{11}$ | $1.5 \times 10^{14}$            | $4.8 \times 10^{14}$            |
| SCSO J052608–561114 | 0.14                 | 0.15             | $41.6 \pm 6.5$   | $63.4 \pm 8.0$         | $1.4 \times 10^{12} \pm 2.6 \times 10^{11}$ | $3.5 \times 10^{14}$            | $3.2 \times 10^{14}$            |
| SCSO J052803–525945 | 0.68                 | 0.69             | $61.5 \pm 7.8$   | $85.3 \pm 9.2$         | $4.3 \times 10^{12} \pm 4.4 \times 10^{11}$ | $6.8 \times 10^{14}$            | $1.2 \times 10^{15}$            |
| SCSO J052810–514839 | 0.65                 | 0.64             | $24.6 \pm 5.0$   | $23.6 \pm 4.8$         | $1.7 \times 10^{12} \pm 1.3 \times 10^{11}$ | $2.3 \times 10^{14}$            | $5.0 \times 10^{14}$            |
| SCSO J052858–535744 | 0.70                 | 0.70             | $37.4 \pm 6.1$   | $53.6 \pm 7.3$         | $3.0 \times 10^{12} \pm 3.1 \times 10^{11}$ | $4.2 \times 10^{14}$            | $8.1 \times 10^{14}$            |
| SCSO J052951–551611 | 0.66                 | 0.65             | $26.7 \pm 5.2$   | $30.3 \pm 5.5$         | $2.4 \times 10^{12} \pm 1.9 \times 10^{11}$ | $3.3 \times 10^{14}$            | $8.1 \times 10^{14}$            |
| SCSO J053052–552056 | 0.73                 | 0.71             | $40.8 \pm 6.4$   | $71.3 \pm 8.4$         | $4.4 \times 10^{12} \pm 4.2 \times 10^{11}$ | $5.8 \times 10^{14}$            | $1.3 \times 10^{15}$            |
| SCSO J053154–552031 | 0.23                 | 0.21             | $92.3 \pm 9.6$   | $114.5 \pm 10.7$       | $2.6 \times 10^{12} \pm 5.1 \times 10^{11}$ | $7.9 \times 10^{14}$            | $6.6 \times 10^{14}$            |
| SCSO J053327–542016 | 0.23                 | 0.25             | $47.7 \pm 6.9$   | $77.3 \pm 8.8$         | $1.7 \times 10^{12} \pm 3.3 \times 10^{11}$ | $3.3 \times 10^{14}$            | $3.4 \times 10^{14}$            |
| SCSO J053437–552312 | 0.76                 | 0.80             | $15.4 \pm 3.9$   | $8.4 \pm 2.9$          | $1.4 \times 10^{12} \pm 1.3 \times 10^{11}$ | $1.1 \times 10^{14}$            | $5.1 \times 10^{14}$            |
| SCSO J053448–543534 | 0.65                 | 0.65             | $23.0 \pm 4.8$   | $24.0 \pm 4.9$         | $1.3 \times 10^{12} \pm 1.3 \times 10^{11}$ | $2.4 \times 10^{14}$            | $3.9 \times 10^{14}$            |
| SCSO J053500–532018 | 0.59                 | 0.57             | $23.1 \pm 4.8$   | $22.8 \pm 4.8$         | $3.2 \times 10^{12} \pm 1.3 \times 10^{11}$ | $2.9 \times 10^{14}$            | $1.2 \times 10^{15}$            |
| SCSO J053632–553123 | 0.72                 | 0.72             | $54.5 \pm 7.4$   | $98.8 \pm 9.9$         | $4.5 \times 10^{12} \pm 5.7 \times 10^{11}$ | $6.2 \times 10^{14}$            | $1.1 \times 10^{15}$            |
| SCSO J053638–553854 | 0.74                 | 0.71             | $40.7 \pm 6.4$   | $65.9 \pm 8.1$         | $2.2 \times 10^{12} \pm 3.1 \times 10^{11}$ | $3.9 \times 10^{14}$            | $4.8 \times 10^{14}$            |
| SCSO J053645–553302 | 0.74                 | 0.71             | $37.1 \pm 6.1$   | $62.7 \pm 7.9$         | $2.0 \times 10^{12} \pm 3.2 \times 10^{11}$ | $4.4 \times 10^{14}$            | $4.9 \times 10^{14}$            |
| SCSO J053655–553809 | 0.76                 | 0.72             | $41.8 \pm 6.5$   | $64.7 \pm 8.1$         | $2.9 \times 10^{12} \pm 3.2 \times 10^{11}$ | $4.6 \times 10^{14}$            | $7.2 \times 10^{14}$            |
| SCSO J053715–541530 | 0.49                 | 0.51             | $21.1 \pm 4.6$   | $20.1 \pm 4.5$         | $1.1 \times 10^{12} \pm 1.1 \times 10^{11}$ | $2.0 \times 10^{14}$            | $3.1 \times 10^{14}$            |
| SCSO J053732–542521 | 0.62                 | 0.61             | $21.5 \pm 4.6$   | $21.4 \pm 4.6$         | $2.4 \times 10^{12} \pm 1.3 \times 10^{11}$ | $2.5 \times 10^{14}$            | $8.6 \times 10^{14}$            |
| SCSO J053952–561423 | 0.36                 | 0.36             | $36.9 \pm 6.1$   | $48.4 \pm 7.0$         | $1.1 \times 10^{12} \pm 1.4 \times 10^{11}$ | $4.1 \times 10^{14}$            | $2.9 \times 10^{14}$            |
| SCSO J054012–561700 | 0.38                 | 0.38             | $51.4 \pm 7.2$   | $59.4 \pm 7.7$         | $1.5 \times 10^{12} \pm 1.8 \times 10^{11}$ | $5.0 \times 10^{14}$            | $3.9 \times 10^{14}$            |
| SCSO J054022–541622 | 0.51                 | 0.48             | $35.1 \pm 5.9$   | $46.1 \pm 6.8$         | $1.8 \times 10^{12} \pm 1.9 \times 10^{11}$ | $3.3 \times 10^{14}$            | $4.3 \times 10^{14}$            |
| SCSO J054052–551943 | 0.76                 | 0.78             | $40.9 \pm 6.4$   | $57.5 \pm 7.6$         | $3.0 \times 10^{12} \pm 3.2 \times 10^{11}$ | $4.5 \times 10^{14}$            | $8.1 \times 10^{14}$            |
| SCSO J054228–525002 | 0.65                 | 0.66             | $21.8 \pm 4.7$   | $21.3 \pm 4.6$         | $1.7 \times 10^{12} \pm 1.3 \times 10^{11}$ | $1.8 \times 10^{14}$            | $4.5 \times 10^{14}$            |
| SCSO J054332–505651 | 0.35                 | 0.36             | $45.5 \pm 6.8$   | $64.0 \pm 8.0$         | $1.3 \times 10^{12} \pm 1.9 \times 10^{11}$ | $4.1 \times 10^{14}$            | $2.8 \times 10^{14}$            |
| SCSO J054358–531349 | 0.24                 | 0.25             | $30.0 \pm 5.5$   | $41.2 \pm 6.4$         | $9.9 \times 10^{11} \pm 1.8 \times 10^{11}$ | $3.1 \times 10^{14}$            | $2.3 \times 10^{14}$            |
| SCSO J054401–511254 | 0.28                 | 0.28             | $31.4 \pm 5.6$   | $43.3 \pm 6.6$         | $1.2 \times 10^{12} \pm 1.6 \times 10^{11}$ | $3.4 \times 10^{14}$            | $3.0 \times 10^{14}$            |
| SCSO J054407–530924 | 0.25                 | 0.26             | $42.9 \pm 6.5$   | $47.4 \pm 6.9$         | $1.0 \times 10^{12} \pm 1.9 \times 10^{11}$ | $3.3 \times 10^{14}$            | $2.2 \times 10^{14}$            |
| SCSO J054436–550319 | 0.35                 | 0.36             | $33.0 \pm 5.8$   | $41.5 \pm 6.4$         | $1.9 \times 10^{12} \pm 1.6 \times 10^{11}$ | $3.6 \times 10^{14}$            | $5.3 \times 10^{14}$            |
| SCSO J054721–554906 | 0.59                 | 0.59             | $38.1 \pm 6.2$   | $49.2 \pm 7.0$         | $2.4 \times 10^{12} \pm 2.1 \times 10^{11}$ | $4.2 \times 10^{14}$            | $6.8 \times 10^{14}$            |
| SCSO J054742–554836 | 0.50                 | 0.50             | $26.6 \pm 5.2$   | $25.8 \pm 5.1$         | $1.3 \times 10^{12} \pm 1.2 \times 10^{11}$ | $2.5 \times 10^{14}$            | $3.9 \times 10^{14}$            |
| SCSO J054811–555601 | 0.64                 | 0.64             | $27.4 \pm 5.2$   | $28.7 \pm 5.3$         | $2.2 \times 10^{12} \pm 1.5 \times 10^{11}$ | $2.8 \times 10^{14}$            | $6.6 \times 10^{14}$            |
| SCSO J054931–522655 | 0.38                 | 0.39             | $21.2 \pm 4.6$   | $22.0 \pm 4.7$         | $2.1 \times 10^{12} \pm 1.1 \times 10^{11}$ | $2.7 \times 10^{14}$            | $7.5 \times 10^{14}$            |
| SCSO J054949–513503 | 0.28                 | 0.28             | $38.4 \pm 6.2$   | $51.4 \pm 7.2$         | $1.3 \times 10^{12} \pm 1.8 \times 10^{11}$ | $4.2 \times 10^{14}$            | $3.3 \times 10^{14}$            |
| SCSO J055017–534601 | 0.49                 | 0.47             | $30.8 \pm 5.5$   | $29.9 \pm 5.5$         | $1.4 \times 10^{12} \pm 1.4 \times 10^{11}$ | $2.5 \times 10^{14}$            | $3.6 \times 10^{14}$            |


**Note.** — Catalog with the optical properties of clusters with mass estimates  $> 3 \times 10^{14} M_{\odot}$  in the 5hr region. For each cluster we note the BCGs photometric redshift and the median photometric redshift for the clusters using the members within 250 kpc of the center of the cluster. The ID is based on the position of the BCG.

that study which we do not recover. One of these (SPT-CL 0509–5342) falls below the mass threshold used here with mass estimates of  $2.1 \times 10^{14} M_\odot$  and  $1.2 \times 10^{14} M_\odot$  for  $M(N_{200})$  and  $M(L_{200})$  respectively. These are lower than the optical mass quoted in Menanteau & Hughes (2009) due to the change in the Reyes et al. (2008) mass parametrization but still are comparable to the weak lensing mass range,  $1.9 - 5.6 \times 10^{14} M_\odot$ , obtained by McInnes et al. (2009) using the same optical dataset. The other cluster (SPT-CL0547–5345) has a photometric redshift of 0.88 (Menanteau & Hughes 2009) that puts it beyond the redshift threshold ( $z = 0.8$ ) we use here.

#### 3.4. *ROSAT* archival data

We searched for X-ray counterparts to the SCS optical clusters using the *ROSAT* All Sky Survey (RASS) data following the same procedure as in Menanteau & Hughes (2009). The raw X-ray photon event lists and exposure maps were downloaded from the MPE *ROSAT* Archive<sup>5</sup> and queried with our own custom software. At the position of each SCS cluster, RASS count rates in the 0.5–2 keV band (corresponding to PI channels 52–201) were extracted from within a  $3'$  radius for the source emission and from within a surrounding annulus ( $5'$  to  $25'$  inner and outer radii) for the background emission. The background-subtracted count rates were converted to X-ray luminosity (in the 0.5–2.0 keV band) assuming a thermal spectrum ( $kT = 5$  keV) and the Galactic column density of neutral hydrogen appropriate to the source position, using data from the Leiden/Argentine/HI Bonn survey (Kalberla et al. 2005). X-ray masses within an overdensity of 500 times the critical density were then determined using the  $L_X$  vs.  $M_{500}$  scaling relation from Vikhlinin et al. (2009). These were then converted to an overdensity of 200 with respect to the average density of the Universe for comparison to the optically-derived masses using a simple multiplicative scaling factor of 1.77. This factor is good to 10% over the redshift and mass range of our sample. In Table 6 we show redshifts, column densities, rates, luminosities, and masses only for those clusters with X-ray emission significant at  $2\sigma$  or higher, while Figure 6 shows all the SCS clusters with either actual values for the X-ray mass or (for most cases) upper limits. We also show points corresponding to a simple stacking of the X-ray and the optical masses for clusters in three redshift and three optical mass bins as the green and red dots, respectively. Each point represents an average of  $\sim 20$  clusters each with some positive X-ray count rate, excluding the several higher significance individual cases plotted in the Figure. It is encouraging that the points scatter by about a factor of two around the line of equal optical and X-ray masses, suggesting that the X-ray emission is in fact associated with hot gas in the potential well of a massive cluster. In at least two cases, we have recovered known clusters (SCSO J051637–543001 and SCSO J232653–524149). And two of the RASS-correlated clusters (SCSO J053154–552031 and SCSO J233227–535827) are among those shown in the color images (i.e., see the bottom left panel of Fig. 3 and the top panel of Fig. 4).

#### 3.5. *XMM-Newton* archival data



**Figure 6.** Plot of X-ray- vs. optically-derived masses for all SCS clusters. The optical mass is the average of  $M(N_{200})$  and  $M(L_{200})$ , while the X-ray mass comes from the X-ray luminosity assuming a  $M-L_X$  scaling relation. Clusters with X-ray emission are shown as the box symbols, while the others are shown as upper limits. The solid line denotes equality; the dashed lines indicate a factor of two range in mass. The green points show the average optical and X-ray masses for clusters stacked in three redshift bins (0.0–0.4, 0.4–0.6, 0.6–0.8) ordered from bottom to top, while the red points were stacked in three optical mass bins ( $<3.2 \times 10^{14} M_\odot$ ,  $3.2 \times 10^{14} M_\odot$ – $4.5 \times 10^{14} M_\odot$ , and  $>4.5 \times 10^{14} M_\odot$ ) ordered from left to right.

There are a number of *XMM-Newton* pointings that overlap with the SCS especially in the 23-hr region. Table 7 lists all the optical clusters from this study (Tables 3 and 4) that are located within the field of view of an archival *XMM-Newton* pointing (specifically the following ObsIds: 0205330301, 0505380601, 0505381801, 0505382201, and 0505383601). In several cases, there is an associated X-ray source detected by the Pipeline Processing System (PPS) which is run on all *XMM-Newton* pointings to produce standard results and are provided to the observer. If a PPS source exists, we list its ID, the offset in arcseconds between the X-ray source and the optical cluster position and whether or not the PPS has flagged the X-ray source as being extended. To determine fluxes and luminosities, we extract spectra from circular regions chosen to maximize the extracted count rate for the extended X-ray sources. For the other sources we use a  $1'$  radius circle. Background spectra come from a surrounding annular region with sufficient area to obtain good photon statistics. The rate, summed over the single PN and both MOS detectors, is quoted in column 7 of Table 7. We also used standard *XMM-Newton* software tools to calculate the instrumental response functions (i.e., the *arf* and *rmf* files) for each cluster. X-ray luminosity values were determined using the extracted spectra and response files assuming a simple source emission model consisting of an absorbed thermal plasma model. For the absorption component we fixed the column density of neutral hydrogen to the Galactic value toward the cluster position (obtained in the same

<sup>5</sup> <ftp://ftp.xray.mpe.mpg.de/rosat/archive/>

way as for the RASS discussed above). Three clusters have enough signal that their mean temperatures can be measured (see below); for the others we assume  $kT = 5$  keV for the luminosity calculation. The final column lists the cluster mass inferred from the X-ray luminosity as discussed above.

The *XMM-Newton* spectra constrain the cluster temperatures for SCSO J051558–543906 and SCSO J231651–545356 to values of  $kT = 1.8^{+0.5}_{-0.3}$  keV and  $kT = 3.7^{+0.6}_{-0.5}$  keV, respectively (errors are at the 68% confidence interval), assuming a metal abundance of 0.3 times the solar value. SCSO J051637–543001 (Abell S0520) is a bright cluster that was the target of a specific *XMM-Newton* program. Our analysis finds a best fit temperature of  $kT = 7.7 \pm 0.3$  keV and metal abundance of  $0.17 \pm 0.04$  relative to solar, which are both consistent with previous work (e.g., Zhang et al. 2006). The X-ray masses for these three systems just mentioned agree quite well with the optically-derived masses and, in the case of SCSO J051637–543001, with the ROSAT-derived mass as well. For the three other clusters in the 23hr region in Table 7 the X-ray masses are all lower than the optical masses by factors of 3–5. For the remaining system (SCSO J051613–542620) there is a catastrophic disagreement between the inferred optical and X-ray masses with the X-ray mass being more than an order of magnitude below the optical one. This is likely due to significant contamination of the optical number counts by cluster members from the nearby, rich system Abell S0520 which is also close in redshift (0.2952 vs. 0.36).

#### 4. SUMMARY

We have fully processed using an independent custom-built pipeline  $\sim 1$  TB of archival *griz* imaging data from the CTIO Blanco 4-m telescope acquired under the NOAO Large Survey program (05B-0043, PI: Joe Mohr), as part of our own Southern Cosmology Survey. This data volume corresponds to 45 nights of observing over 3 years (2005–2007) and covers  $70 \text{ deg}^2$  of the southern sky that has also been fully observed by ACT and SPT. Here we have presented the first results from the full nominal data set, namely a sample of 105 massive, optically-selected galaxy clusters. Future studies will present the properties of lower mass clusters and groups as well as multi-wavelength studies of cluster physics utilizing selected clusters from this sample.

The current sample is limited to systems with optically-derived masses greater than  $3 \times 10^{14} M_{\odot}$  and redshifts less than 0.8. We have chosen this mass limit to be at or below the anticipated mass threshold of ACT and SPT in order to encompass the upcoming significant SZE detections. However we also have a redshift limit which is due to the depth of the imaging and the wavelength coverage of the filter set. Thus we are missing the most interesting massive clusters at high redshift ( $z > 0.8$ ). However, as demonstrated in Menanteau & Hughes (2009), the optical data analyzed here can be used to confirm the presence of a cluster when conducting a targeted positional search for a high significance SZE candidate.

The recent success of the mm-band wide-survey-area experiments (ACT and SPT) in finding new clusters

through untargeted SZE surveys has been a strong catalyst for our work. We present this cluster sample to aid in the verification of SZE cluster candidates and the characterization of the SZE selection function which currently is observationally unexplored. Moreover, we anticipate stacking the ACT mm-band maps at the positions of optical clusters to detect, statistically, the average SZE signal for systems that fall below the ACT detection threshold for individual sources.

Given the large volume of this data set we believe it might be helpful to address other problems in astrophysics; therefore we plan to make the Blanco data products (i.e., photometric source catalogs and images) available to the community in December 2010 at the following URL <http://scs.rutgers.edu>.

We would like to thank the Blanco Cosmology Survey team for the planning and execution of the CTIO Blanco observations that were used in this paper. We have made use of the ROSAT Data Archive of the Max-Planck-Institut für extraterrestrische Physik (MPE) at Garching, Germany as well as results obtained from the High Energy Astrophysics Science Archive Research Center, provided by NASA’s Goddard Space Flight Center. This research has made use of data obtained from or software provided by the US National Virtual Observatory, which is sponsored by the National Science Foundation. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. Partial financial support was provided by the National Science Foundation under the PIRE program (award number OISE-0530095). We also acknowledge support from NASA/XMM grants NNX08AX55G and NNX08AX72G.

#### REFERENCES

- Annis, J., et al. 1999, Bulletin of the American Astronomical Society, 31, 1391
- Abell, G. O. 1958, ApJS, 3, 211
- Abell, G. O., Corwin, H. G., Jr., & Olowin, R. P. 1989, ApJS, 70, 1
- Benítez, N. 2000, ApJ, 536, 571
- Becker, M. R., et al. 2007, ApJ, 669, 905
- Bertin, E., & Arnouts, S. 1996, A&AS, 117, 393
- Bertin, E., SWarp Resample and Coadd Software, <http://terapix.iap.fr/cpld/oldSite/soft/swarp/index.html>
- Blakeslee, J. P., et al. 2003, ApJ, 596, L143
- Blanton, M. R., et al. 2003, ApJ, 592, 819
- Bower, R. G., Lucey, J. R., & Ellis, R. S. 1992, MNRAS, 254, 601
- Brown, M. J. I., Dey, A., Jannuzi, B. T., Brand, K., Benson, A. J., Brodwin, M., Croton, D. J., & Eisenhardt, P. R. 2007, ApJ, 654, 858
- Dong, F., Pierpaoli, E., Gunn, J. E., & Wechsler, R. H. 2008, ApJ, 676, 868
- Fowler, J. W., et al. 2009, arXiv:1001.2934
- Gal, R. R., de Carvalho, R. R., Odewahn, S. C., Djorgovski, S. G., & Margoniner, V. E. 2000, AJ, 119, 12
- Gal, R. R. 2008, A Pan-Chromatic View of Clusters of Galaxies and the Large-Scale Structure, 740, 119
- Gal, R. R., Lopes, P. A. A., de Carvalho, R. R., Kohl-Moreira, J. L., Capelato, H. V., & Djorgovski, S. G. 2009, AJ, 137, 2981
- Gladders, M. D., & Yee, H. K. C. 2000, AJ, 120, 2148
- Gladders, M. D., Hoekstra, H., Yee, H. K. C., Hall, P. B., & Barrientos, L. F. 2003, ApJ, 593, 48
- Gladders, M. D., & Yee, H. K. C. 2005, ApJS, 157, 1



**Table 5**  
Catalogued Sources Associated with SCS Optical Clusters

| SCS ID              | Catalog name                        | Distance to BCG | Source type             |
|---------------------|-------------------------------------|-----------------|-------------------------|
| SCSO J051412–514004 | SUMSS J051411–513953                | 13''            | RadioS                  |
| SCSO J051637–543001 | 2MASX J05163736–5430017             | 3''             | Galaxy                  |
| "                   | Abell S0520, RXC J0516.6–5430       | 2.4'            | GClstr ( $z = 0.2952$ ) |
| "                   | ACT-CL J0516–5432, SPT-CL 0517–5430 | 2.8', 0.5'      | SZ-GClstr               |
| SCSO J052113–510418 | SUMSS J052114–510419                | 13''            | RadioS                  |
| SCSO J052533–551818 | Abell S0529                         | 1.5'            | GClstr                  |
| SCSO J052608–561114 | Abell S0530                         | 2.4'            | GClstr                  |
| SCSO J052803–525945 | SUMSS J052805–525953                | 24''            | RadioS                  |
| "                   | SPT-CL 0528–5300                    | 0.4'            | SZ-GClstr               |
| SCSO J053327–542016 | 2MASX J05332723–5420154             | 2''             | Galaxy                  |
| SCSO J053437–552312 | SUMSS J053442–552248                | 55''            | RadioS                  |
| SCSO J053632–553123 | SUMSS J053629–553147                | 30''            | RadioS                  |
| SCSO J053715–541530 | SUMSS J053718–541608                | 48''            | RadioS                  |
| SCSO J054012–561700 | SUMSS J054014–561723                | 31''            | RadioS                  |
| SCSO J054407–530924 | SUMSS J054406–530922                | 8''             | RadioS                  |
| SCSO J054949–513503 | SUMSS J054948–513454                | 9''             | RadioS                  |
| SCSO J055017–534601 | SUMSS J055019–534601                | 25''            | RadioS                  |
| SCSO J231455–555308 | 2MASX J23145553–5553093             | 5''             | Galaxy                  |
| SCSO J232001–565222 | SUMSS J232001–565219                | 4''             | RadioS                  |
| SCSO J232653–524149 | RXC J2326.7–5242                    | 1.2'            | GClstr ( $z = 0.1074$ ) |
| "                   | SUMSS J232654–524153                | 14''            | RadioS                  |
| SCSO J233227–535827 | 1RXS J233224.3–535840               | 27''            | XrayS                   |
| SCSO J233544–535115 | SUMSS J233544–535113                | 3''             | RadioS                  |
| SCSO J234156–530848 | SUMSS J234156–530849                | 5''             | RadioS                  |
| SCSO J234703–535051 | SUMSS J234703–535052                | 3''             | RadioS                  |
| SCSO J234917–545521 | SUMSS J234917–545518                | 8''             | RadioS                  |
| SCSO J235055–530124 | SUMSS J235054–530141                | 17''            | RadioS                  |
| SCSO J235138–545253 | 2MASX J23513813–5452538             | 2''             | Galaxy                  |
| "                   | SUMSS J235138–545255                | 3''             | RadioS                  |

**Note.** — Catalogued sources from the NASA/IPAC Extragalactic Database (NED) correlated with SCS optical clusters.

**Table 6**  
Optical Clusters with X-ray counterparts from the ROSAT All Sky Survey

| SCS ID              | $z_{\text{cluster}}$ | $N_{\text{H}}$<br>( $10^{20} \text{ cm}^{-2}$ ) | Rate<br>(cts s $^{-1}$ ) | $L_X(0.5\text{--}2.0 \text{ keV})$<br>( $10^{44} \text{ erg s}^{-1}$ ) | $M_{200}(L_X)$<br>( $M_{\odot}$ ) |
|---------------------|----------------------|-------------------------------------------------|--------------------------|------------------------------------------------------------------------|-----------------------------------|
| SCSO J051136–561045 | 0.70                 | 1.61                                            | $0.015 \pm 0.006$        | $3.0 \pm 1.3$                                                          | $7.9 \times 10^{14}$              |
| SCSO J051637–543001 | 0.2952*              | 2.05                                            | $0.205 \pm 0.030$        | $6.1 \pm 0.9$                                                          | $1.6 \times 10^{15}$              |
| SCSO J052533–551818 | 0.72                 | 4.08                                            | $0.011 \pm 0.005$        | $2.5 \pm 1.2$                                                          | $6.9 \times 10^{14}$              |
| SCSO J053154–552031 | 0.23                 | 5.20                                            | $0.017 \pm 0.006$        | $0.3 \pm 0.1$                                                          | $2.7 \times 10^{14}$              |
| SCSO J053952–561423 | 0.36                 | 4.90                                            | $0.010 \pm 0.005$        | $0.5 \pm 0.3$                                                          | $3.3 \times 10^{14}$              |
| SCSO J054407–530924 | 0.25                 | 5.01                                            | $0.020 \pm 0.005$        | $0.5 \pm 0.1$                                                          | $3.3 \times 10^{14}$              |
| SCSO J054721–554906 | 0.59                 | 6.98                                            | $0.007 \pm 0.004$        | $1.2 \pm 0.6$                                                          | $4.8 \times 10^{14}$              |
| SCSO J054949–513503 | 0.28                 | 4.58                                            | $0.012 \pm 0.004$        | $0.3 \pm 0.1$                                                          | $2.7 \times 10^{14}$              |
| SCSO J232653–524149 | 0.1074*              | 1.28                                            | $0.166 \pm 0.035$        | $0.6 \pm 0.1$                                                          | $4.1 \times 10^{14}$              |
| SCSO J233227–535827 | 0.35                 | 1.28                                            | $0.091 \pm 0.026$        | $3.9 \pm 1.1$                                                          | $1.2 \times 10^{15}$              |

**Note.** — Catalog of the optical clusters associated with X-ray emission in the ROSAT All Sky Survey. The quoted redshifts are just repeated from Tables 3 and 4 except for the starred values which are spectroscopic redshifts from NED (see Table 5). Rates were extracted from within 3' radii circles except for the three brightest clusters, SCSO J051637–543001, SCSO J232653–524149, and SCSO J233227–535827, for which radii of 10', 7', and 6', respectively, were used.

Goto, T., et al. 2002, *AJ*, 123, 1807  
Hansen, S. M., McKay, T. A., Wechsler, R. H., Annis, J., Sheldon, E. S., & Kimball, A. 2005, *ApJ*, 633, 122  
Hincks, A. D., et al. 2009, *arXiv:0907.0461*  
Johnston, D. E., et al. 2007, *ArXiv e-prints*, 709, *arXiv:0709.1159*  
Jenkins, A., Frenk, C. S., White, S. D. M., Colberg, J. M., Cole, S., Evrard, A. E., Couchman, H. M. P., & Yoshida, N. 2001, *MNRAS*, 321, 372  
Kalberla, P. M. W., Burton, W. B., Hartmann, Dap, Arnal, E. M., Bajaja, E., Morras, R., & Pöppel, W. G. L. 2005, *A&A*, 440, 775  
Koester, B. P., et al. 2007, *ApJ*, 660, 221

Koester, B. P., et al. 2007, *ApJ*, 660, 239  
Komatsu, E., et al. 2009, *ApJS*, 180, 330  
Komatsu, E., et al. 2010, *arXiv:1001.4538*  
Kravtsov, A. V., Vikhlinin, A., & Nagai, D. 2006, *ApJ*, 650, 128  
Loh, Y.-S., & Strauss, M. A. 2006, *MNRAS*, 366, 373  
Lopes, P. A. A., de Carvalho, R. R., Gal, R. R., Djorgovski, S. G., Odewahn, S. C., Mahabal, A. A., & Brunner, R. J. 2004, *AJ*, 128, 1017  
Lueker, M., et al. 2009, *arXiv:0912.4317*  
Mauch, T., Murphy, T., Buttery, H. J., Curran, J., Hunstead, R. W., Piestrzynski, B., Robertson, J. G., & Sadler, E. M. 2003, *MNRAS*, 342, 1117

**Table 7**  
Optical Clusters Located within the Field of View of XMM-Newton Observations

| SCS ID              | $z_{\text{cluster}}$ | $N_{\text{H}}$<br>( $10^{20} \text{ cm}^{-2}$ ) | X-ray ID           | Offset<br>( $''$ ) | Extended | Extr Rad<br>(kpc) | Rate<br>( $10^{-3} \text{ cts s}^{-1}$ ) | $L_X$ (0.5–2 keV)<br>( $10^{44} \text{ ergs s}^{-1}$ ) | $M(L_X)$<br>( $10^{14} M_{\odot}$ ) |
|---------------------|----------------------|-------------------------------------------------|--------------------|--------------------|----------|-------------------|------------------------------------------|--------------------------------------------------------|-------------------------------------|
| SCSO J051558–543906 | 0.64                 | 2.07                                            | XMM J051600–543900 | 19                 | yes      | 910 (2.2')        | $15.7 \pm 2.6$                           | 0.60                                                   | 3.0                                 |
| SCSO J051613–542620 | 0.36                 | 1.98                                            | ...                | ...                | ...      | 300 (1')          | $6.8 \pm 1.4$                            | 0.027                                                  | 0.5                                 |
| SCSO J051637–543001 | 0.2952*              | 2.05                                            | XMM J051635–543022 | 25                 | yes      | 1390 (5.3')       | $1740 \pm 15$                            | 5.1                                                    | 14.4                                |
| SCSO J231651–545356 | 0.36                 | 1.29                                            | XMM J231653–545410 | 26                 | yes      | 650 (2.2')        | $73.7 \pm 3.5$                           | 0.71                                                   | 4.1                                 |
| SCSO J232856–552428 | 0.57                 | 1.29                                            | XMM J232856–552429 | 5                  | no       | 390 (1')          | $5.4 \pm 1.3$                            | 0.07                                                   | 0.8                                 |
| SCSO J233420–542732 | 0.55                 | 1.27                                            | ...                | ...                | ...      | 380 (1')          | $5.2 \pm 1.4$                            | 0.08                                                   | 0.9                                 |
| SCSO J233556–560602 | 0.63                 | 1.27                                            | ...                | ...                | ...      | 410 (1')          | $6.2 \pm 1.7$                            | 0.10                                                   | 1.0                                 |

**Note.** — Luminosities calculated by fitting an absorbed thermal emission model with a fixed 5 keV temperature, except for SCSO J051558–543906, SCSO J051637–543001, and SCSO J231651–545356 for which there was enough signal to determine their temperatures (see text). The quoted redshifts are just repeated from Tables 3 and 4 except for the starred value which is a spectroscopic redshift from NED (see Table 5).

- McInnes, R. N., Menanteau, F., Heavens, A. F., Hughes, J. P., Jimenez, R., Massey, R., Simon, P., & Taylor, A. 2009, MNRAS, 399, L84
- Menanteau, F., et al. 2009, ApJ, 698, 1221
- Menanteau, F., & Hughes, J. P. 2009, ApJ, 694, L136
- Okabe, N., Takada, M., Umetsu, K., Futamase, T., & Smith, G. P. 2010, PASJ, 62, 811
- Postman, M., Lubin, L. M., Gunn, J. E., Oke, J. B., Hoessel, J. G., Schneider, D. P., & Christensen, J. A. 1996, AJ, 111, 615
- Ramella, M., Boschini, W., Fadda, D., & Nonino, M. 2001, A&A, 368, 776
- Reiprich, T. H., & Böhringer, H. 2002, ApJ, 567, 716
- Reyes, R., Mandelbaum, R., Hirata, C., Bahcall, N., & Seljak, U. 2008, MNRAS, 390, 1157
- Rozo, E., et al. 2009, ApJ, 703, 601
- Rozo, E., et al. 2009, ApJ, 699, 768
- Rozo, E., et al. 2010, ApJ, 708, 645
- Rykoff, E. S., et al. 2008, MNRAS, 387, L28
- Schechter, P. 1976, ApJ, 203, 297
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, ApJ, 500, 525
- Skrutskie, M. F., et al. 2006, AJ, 131, 1163
- Sheldon, E. S., et al. 2009, ApJ, 703, 2217
- Smith, J. A., Allam, S. S., Tucker, D. L., Stute, J. L., Rodgers, C. T., & Stoughton, C. 2007, AJ, submitted
- Sunyaev, R. A., & Zeldovich, Y. B. 1972, Comments on Astrophysics and Space Physics, 4, 173
- Staniszewski, Z., et al. 2009, ApJ, 701, 32
- Valdes, F. G. 1998, Astronomical Data Analysis Software and Systems VII, 145, 53
- Vikhlinin, A., et al. 2009, ApJ, 692, 1033
- Voges, W., et al. 1999, A&A, 349, 389
- Yasuda, N., et al. 2001, AJ, 122, 1104
- Zhang, Y.-Y., Böhringer, H., Finoguenov, A., Ikebe, Y., Matsushita, K., Schuecker, P., Guzzo, L., & Collins, C. A. 2006, A&A, 456, 55